

Collective excitations of Bose-Einstein condensed gases at finite temperatures

R. J. Dodd*

Institute for Physical Science and Technology, University of Maryland at College Park, College Park, MD 20742.

Mark Edwards*

Department of Physics, Georgia Southern University, Statesboro, GA 30460-8031.

Charles W. Clark

Electron and Optical Physics Division, Physics Laboratory, National Institute of Standards and Technology, Technology Administration, U. S. Department of Commerce, Gaithersburg, MD 20899-0001.

K. Burnett*

Clarendon Laboratory, Department of Physics, University of Oxford, Parks Road, Oxford OX1 3PU, United Kingdom.

(February 7, 2008)

We have applied the Popov version of the Hartree-Fock-Bogoliubov (HFB) approximation to calculate the finite-temperature excitation spectrum of a Bose-Einstein condensate (BEC) of ^{87}Rb atoms. For lower values of the temperature, we find excellent agreement with recently-published experimental data for the JILA TOP trap. In contrast to recent comparison of the results of HFB-Popov theory with experimental condensate fractions and specific heats, there is disagreement of the theoretical and recent experimental results near the BEC phase transition temperature.

PACS Numbers: 3.75.Fi, 67.40.Db, 67.90.+Z

Laboratory realizations of gaseous Bose-Einstein condensates (BECs) [1–3] have prompted vigorous experimental [4] investigations of the temperature-dependent properties of these mesoscopic quantum systems. The current theoretical interest in such condensates derives, in part, from the fact that experimental tests of many-body theories that are thought to apply to BEC — a phenomenon occurring in many areas of physics — can be performed for the first time. An accurate theory of such systems is therefore of fundamental interest, and will also have practical applications. In this Letter, we explore the limits of validity of the simplest temperature-dependent mean-field theory — a simplified version of the Hartree-Fock-Bogoliubov (HFB) approximation originally introduced by Popov [5] — by presenting the first comparison with experiment of this theory's predictions for condensate excitation spectra at $T > 0$.

Condensate properties predicted by such theories include condensate and thermal-atom spatial density profiles, condensate fractions, specific heats, and excitation frequencies. Of these properties, excitation spectra provide the most sensitive test of the applicability of competing theories since the other quantities listed depend on sums over states and are thus insensitive to small errors in the excitation spectrum. One example of this can be found in the approach of zero-temperature exci-

tation frequencies to the Thomas-Fermi limit (i.e., the limit $N_0 \rightarrow \infty$, where N_0 is the number of condensate atoms) as N_0 increases. For ^{87}Rb condensates confined in the JILA TOP (time-averaged orbiting potential) trap where $N_0 > 4000$ atoms, the Thomas-Fermi predictions for spatial density profiles are extremely accurate while low-order excitation frequencies can differ by as much as 10%.

There is good *a priori* reason to expect that the HFB-Popov theory should provide good predictions of experimental T -dependent collective excitation frequencies. Measurements of zero-temperature frequencies [6] exhibited excellent agreement with the predictions of zero-temperature, mean field theory [7]. Furthermore, semiclassical variants of the HFB-Popov theory have exhibited excellent agreement with experiment for T -dependent condensate fractions and specific heats for temperatures up to near T_c [8]. The HFB-Popov theory is a finite-temperature extension of mean-field theory which provides self-consistent treatment of the condensed and thermal components of the gas and which should describe the linear response of the condensate to small-amplitude mechanical disturbances [9].

Although the HFB-Popov equations have been derived elsewhere [9,10], we shall briefly state the physics behind the basic equations here. The confined Bose gas is portrayed as a thermodynamic equilibrium system under the grand canonical ensemble whose thermodynamic variables are N , the total number of trapped atoms, T , the absolute temperature, and either N_0 or μ , the chemical potential. The system hamiltonian has the form

$$K \equiv H - \mu N = \int d\mathbf{r} \hat{\psi}^\dagger(\mathbf{r})(H_0 - \mu)\hat{\psi}(\mathbf{r}) + \frac{U_0}{2} \int d\mathbf{r} \hat{\psi}^\dagger(\mathbf{r})\hat{\psi}^\dagger(\mathbf{r})\hat{\psi}(\mathbf{r})\hat{\psi}(\mathbf{r}) . \quad (1)$$

where $\hat{\psi}(\mathbf{r})$ is the Bose field operator which annihilates an atom at position $\mathbf{r}$ and $H_0 = \frac{\hbar^2}{2M} \nabla^2 + V_{\text{trap}}(\mathbf{r})$ is the bare trap Hamiltonian. For the system treated here,

the trap potential $V_{\text{trap}}(\mathbf{r}) = M(\omega_\rho^2 \rho^2 + \omega_z^2 z^2)/2$, where M is the atomic mass, and ω_ρ and ω_z are the radial and axial trap frequencies, respectively. The quantity $U_0 = 4\pi\hbar^2 a/M$ is a measure of the interaction strength between atoms, with a being the scattering length for zero-energy binary atomic collisions, taken to be $109a_0$ for ^{87}Rb [11], where a_0 is the Bohr radius.

The Bose field operator is decomposed into a c -number condensate wave function plus an operator describing the non-condensate part: $\hat{\psi}(\mathbf{r}) = N_0^{1/2}\phi(\mathbf{r}) + \tilde{\psi}(\mathbf{r})$ and inserted into Eq. (1). When terms cubic and quartic in $\tilde{\psi}(\mathbf{r})$ are treated within the mean-field approximation the grand-canonical hamiltonian reduces to a sum of four terms: $K = K_0 + K_1 + K_1^\dagger + K_2$. The first term, K_0 is a c -number, the second and third terms are linear in $\tilde{\psi}(\mathbf{r})$ and $\tilde{\psi}^\dagger(\mathbf{r})$ and the last term is quadratic in these quantities. It is easy to show that the linear terms vanishes identically if $\phi(\mathbf{r})$ satisfies the generalized Gross-Pitaevskii (GP) equation

$$\left\{ H_0 + U_0 \left[N_0 |\phi(\mathbf{r})|^2 + 2\tilde{n}(\mathbf{r}) \right] \right\} \phi(\mathbf{r}) = \mu \phi(\mathbf{r}), \quad (2)$$

where $\tilde{n}(\mathbf{r})$ is the density of non-condensate atoms. Note that the condensate wavefunction is normalized to unity.

The term K_2 has the form

$$K_2 = \int d\mathbf{r} \tilde{\psi}^\dagger(\mathbf{r}) \mathcal{L} \tilde{\psi}(\mathbf{r}) + \frac{N_0 U_0}{2} \int d\mathbf{r} (\phi)^2(\mathbf{r}) \tilde{\psi}^\dagger(\mathbf{r}) \tilde{\psi}(\mathbf{r}) + \frac{N_0 U_0}{2} \int d\mathbf{r} (\phi^*)^2(\mathbf{r}) \tilde{\psi}(\mathbf{r}) \tilde{\psi}^\dagger(\mathbf{r}). \quad (3)$$

where $\mathcal{L} \equiv H_0 + 2U_0 n(\mathbf{r}) - \mu$ and $n(\mathbf{r}) = N_0 |\phi(\mathbf{r})|^2 + \tilde{n}(\mathbf{r})$ is the total trapped-atom density. The term K_2 can be diagonalized by the following Bogoliubov transformation

$$\begin{aligned} \tilde{\psi}(\mathbf{r}) &= \sum_j (u_j(\mathbf{r}) \alpha_j + v_j^*(\mathbf{r}) \alpha_j^\dagger), \\ \tilde{\psi}^\dagger(\mathbf{r}) &= \sum_j (u_j^*(\mathbf{r}) \alpha_j^\dagger + v_j(\mathbf{r}) \alpha_j). \end{aligned} \quad (4)$$

if the quasi-particle amplitudes $u_j(\mathbf{r})$ and $v_j(\mathbf{r})$, satisfy the coupled HFB-Popov equations:

$$\begin{aligned} \mathcal{L} u_j(\mathbf{r}) + N_0 U_0 |\phi(\mathbf{r})|^2 v_j(\mathbf{r}) &= E_j u_j(\mathbf{r}) \\ \mathcal{L} v_j(\mathbf{r}) + N_0 U_0 |\phi(\mathbf{r})|^2 u_j(\mathbf{r}) &= -E_j v_j(\mathbf{r}), \end{aligned} \quad (5)$$

The α_j and $\alpha_j^\dagger$ are quasi-particle annihilation and creation operators that satisfy the usual Bose commutation relations.

The density of the thermal component of the gas $\tilde{n}(\mathbf{r}) \equiv \langle \tilde{\psi}^\dagger \tilde{\psi} \rangle$ and thus can be written in terms of the quasi-particle amplitudes as

$$\tilde{n}(\mathbf{r}) = \sum_j \left\{ \left[|u_j(\mathbf{r})|^2 + |v_j(\mathbf{r})|^2 \right] N_j + |v_j(\mathbf{r})|^2 \right\}, \quad (6)$$

where $N_j = (e^{\beta E_j} - 1)^{-1}$ is the Bose-Einstein factor, and $\beta = (k_B T)^{-1}$ with k_B the Boltzmann constant. The total number of trapped atoms, N , is given by

$$N = \int d\mathbf{r} n(\mathbf{r}) = N_0 + \int d\mathbf{r} \tilde{n}(\mathbf{r}). \quad (7)$$

Our version of Eq. (5) differs from that of Ref. [12] via a sign change in the definition of $v_j(\mathbf{r})$.

Equations (2), (5), (6), and (7) form a closed system of equations that we have referred to as the ‘‘HFB-Popov’’ equations. We have numerically solved these equations under conditions appropriate to ^{87}Rb atoms confined in the JILA TOP trap [13]. We choose our state variables to be $\{T, \mu, N\}$, fix T and μ , and then determine N by solving the HFB-Popov equations. This is equivalent to the alternative triple of state variables $\{T, N_0, N\}$, since there is a one-to-one relationship between N_0 and μ .

We have solved the HFB-Popov equations by an iterative procedure, each cycle of the iteration consisting of two steps. In the first step of each cycle, we solve Eq. (2) for new values of $\phi(\mathbf{r})$ and N_0 with a basis-set approach as described previously [14] using $\tilde{n}(\mathbf{r})$ obtained in the previous cycle. In the second step, we solve Eqs. (5) using $\tilde{n}(\mathbf{r})$ from the previous cycle, and the newly generated values of $\phi(\mathbf{r})$ and N_0 . With the quasi-particle amplitudes expanded in the trap basis, Eqs. (5) yield a generalized matrix eigenvalue problem for the basis-set coefficients. We recast the generalized matrix eigenvalue problem by using a decoupling transformation consisting of taking the sum and difference of Eqs. (5). This transformation is equivalent to that of Hutchinson *et al.* [12], except that it is expressed in terms of basis set expansion coefficients. Completion of this step yields the $\{u_j(\mathbf{r}), v_j(\mathbf{r})\}$, and E_j which are used in Eq. (6) to update $\tilde{n}(\mathbf{r})$. Equation (7) then updates the total number of trapped atoms, N . Convergence is reached when the change in N from one cycle to the next is smaller than a specified tolerance. To obtain converged results at high-temperatures, we add a correction to the total number of atoms, N , at each iteration cycle. High-energy quasi-particle eigenfunctions have negligible overlap with the condensate wave function, so their presence in the thermal sum of Eq.(6) does not significantly modify the low-lying excitation frequencies, but does contribute to the value of N .

We have checked the accuracy of our numerical work by writing two independent codes, which produce identical answers. The ideal gas result is recovered when we set $a = 0$, and we have reproduced the results reported in Ref. [12]. We now discuss the comparison of this approach with experiment.

Figure 1 compares the experimental [13] excitation spectrum of ^{87}Rb in the JILA TOP trap, *vs.* our HFB-Popov results for the $m = 0$ and $m = 2$ modes. The abscissa is the scaled temperature $T' = T/T_0(N, \bar{\omega})$ where

$T_0(N, \omega) \equiv \hbar\omega/k_B [N/\zeta(3)]^{1/3}$ is the theoretical transition temperature for an ideal, trapped Bose gas and $\bar{\omega} = (\omega_\rho^2 \omega_z)^{1/3}$. The ordinate is the excitation frequency expressed in units of ω_ρ . Our results were obtained using the experimental value of T , and a value of μ that yielded the experimentally determined value of N . Thus, as for our previous treatment of zero-temperature excitation spectra [7], *this calculation contains no adjustable parameters*. The agreement between theory and experiment is very good (on the order of 5%) for low and intermediate temperatures ($T' \leq 0.65$). It should be noted that, as depicted in Fig. 2, the high end of this temperature range corresponds to a non-condensate fraction of about 50%. However, as the temperature increases, the HFB-Popov excitation frequencies diverge from the experimental data. This feature of the comparison holds true for both $m = 0$ and $m = 2$ modes.

The behavior of the calculated excitation frequencies can be understood in a simple way. The HFB-Popov equations determine the equation of state for the state variables $\{N, N_0, T\}$, so that, given the values of N and T , a unique N_0 is determined. For fixed N , this relationship generates the condensate fraction N_0/N as a function of T , which is shown in Fig. 2 for the JILA TOP trap with $N = 2000$. One can easily predict the temperature-dependent mode frequencies for the $N = 2000$ system by finding the number of condensate atoms, N_0 , from Fig. 2, and then determining the *zero-temperature* excitation frequency of a condensate with N_0 atoms, which is a much simpler calculation. Figure 3 shows a comparison of the three lowest frequencies numerically determined from the HFB-Popov equations, and by the equivalent $T = 0$ method just described. The agreement of these two approaches is very good over nearly the entire temperature range. The two solid curves in Fig. 1 are the frequencies determined by the same procedure except that the number of condensate atoms was taken from experiment. In short, the principal effect of finite temperature on the HFB-Popov excitation spectra is largely an effect of *condensate depletion*: the dynamics of the finite-temperature condensate are essentially the same as those of a zero-temperature condensate with the same value of N_0 . This is consistent with earlier calculations [15,16] of the speed of sound in a homogeneous Bose-condensed gas, which found that its temperature dependence was effectively a condensate-density dependence. We discuss this result in a broader context in a separate paper [17], in which we show that HFB-Popov results can be reproduced quantitatively by a much simpler “two-gas” model: the condensate gas, which is described by the zero-temperature GP equation; and the thermal cloud, which is described as an ideal Bose gas in an effective potential created by the condensate. This effective potential repels the thermal gas from the condensate gas, which results in the essential independence of tempera-

ture of all condensate properties except N_0 . Application of simple quantum statistical mechanics to this model can generate the full phase diagram of Fig. 2 directly.

As Fig. 1 clearly shows, the HFB-Popov solutions reproduce the experimental results quite well when $T \leq 0.65 T_0$, but fail at higher temperatures. The HFB-Popov formalism is biased toward description of the condensate, as it represents the condensate excitations as taking place in a static thermal cloud. This results in at least one minor failure of the approach, which is weakly visible in Fig. 3 as a deviation of the $m = 1$ mode frequency from unity near T_0 , in violation of the generalized Kohn theorem for parabolic confinement [12,18]. This mode should correspond to a rigid oscillation of the complete N -atom system, and the deviation of its frequency from unity results from the HFB-Popov approximation holding the thermal component fixed and allowing only the condensate to oscillate. In experiments of the type discussed here, however, the thermal component and condensate must both be driven by the modulation of their common confining potential. For other, non-rigid, oscillations we may thus also expect thermal and condensate modes to be coupled in general. Thus, HFB-Popov frequencies will only correspond to the experimental values if the condensate response to mechanical disturbance does not induce modulations of the thermal density, so that there is no back-action of the thermal cloud on the condensate motion. A more general theory that accounts for such condensate-cloud interactions has recently been outlined [19], but remains to be implemented. The self-consistent inclusion of pair terms, which are neglected in the Popov approximation, may also be important for capturing multiple-collision effects in the theory of trapped BECs [20]. The effect of such terms can be very marked close to the transition temperature [21].

In conclusion, we have delineated the region of validity of the Popov version of finite temperature HFB theory by comparing it directly with the results of recent experiments. Good agreement is obtained for condensate fractions from unity down to about 0.5, so the HFB-Popov is apparently correctly describing finite-temperature phenomena in a nontrivial regime. This comparison confirms the critical role of evaporatively cooled gases in establishing proper finite-temperature field theories of Bose-Einstein condensation, and shows that there is still work needed to establish satisfactory agreement between theory and experiment for cases of small condensate fraction.

We thank the JILA group for providing us access to their experimental data. This work was supported in part by the U. S. National Science Foundation under grants PHY-9601261 and PHY-9612728, the U. S. Office of Naval Research, and the U. K. Engineering and Physical Sciences Research Council.

* Also at Physics Laboratory, National Institute of Standards and Technology, Technology Administration, U. S. Department of Commerce, Gaithersburg, MD 20899.

- [1] M.H. Anderson *et al.*, Science **269**, 198 (1995).
- [2] C.C. Bradley, C.A. Sackett, J.J. Tollett, and R.G. Hulet, Phys. Rev. Lett. **75**, 1687 (1995).
- [3] K.B. Davis *et al.*, Phys. Rev. Lett. **75**, 3969 (1995).
- [4] J.R. Ensher *et al.*, Phys. Rev. Lett. **78**, 764 (1997); D.S. Jin *et al.*, Phys. Rev. Lett. **77**, 420 (1996); M.-O. Mewes *et al.*, Phys. Rev. Lett. **77**, 988 (1996).
- [5] V.N. Popov, *Functional Integrals and Collective Modes* (Cambridge University Press, New York, 1987), Ch. 6.
- [6] D.S. Jin *et al.*, Phys. Rev. Lett. **77**, 420 (1996); M.-O. Mewes *et al.*, Phys. Rev. Lett. **77**, 992 (1996).
- [7] M. Edwards *et al.*, Phys. Rev. Lett. **77**, 1671 (1996); S. Stringari, Phys. Rev. Lett. **77**, 2360 (1996).
- [8] A. Minguzzi, S. Conti, and M. P. Tosi, J. Phys.: Condens. Matter, vol. 9, L33 (1997); S. Giorgini, L. P. Pitaevskii, and S. Stringari, Phys. Rev. A **54**, 4633 (1996); S. Giorgini, L.P. Pitaevskii, and S. Stringari, "Thermodynamics of a trapped, Bose-condensed gas", unpublished (cond-mat/9704014).
- [9] A. Griffin, Phys. Rev. B **53**, 9341 (1996).
- [10] A.L. Fetter, Ann. Phys. (NY) **70**, 1671 (1972).
- [11] D. Heinzen, private communication.
- [12] D.A.W. Hutchinson, E. Zaremba, and A. Griffin, Phys. Rev. Lett. **78**, 1842 (1996).
- [13] D.S. Jin *et al.*, Phys. Rev. Lett. **78**, 764 (1997).
- [14] M. Edwards *et al.*, Phys. Rev. A **53**, R1950 (1996); J. Res. Nat. Inst. Stand. Technol. **101**, 553 (1996).
- [15] P. Szépfalusy and I. Kondor, Ann. Phys. (N.Y.) **82**, 1 (1974).
- [16] S. H. Payne and A. Griffin, Phys. Rev. B **32**, 7199 (1985).
- [17] R.J. Dodd, K. Burnett, M. Edwards, and C. W. Clark, Phys. Rev. A (submitted).
- [18] See J.F. Dobson, Phys. Rev. Lett. **73**, 2244 (1994); and references therein.
- [19] N.P. Proukakis and K. Burnett, J. Res. Nat. Inst. Stand. Technol. **101**, 457 (1996).
- [20] N.P. Proukakis, K. Burnett, and H.T.C. Stoof, preprint (1997).
- [21] M. Bijlsma and H.T.C. Stoof, Phys. Rev. A **55**, 498 (1997).




FIG. 1. The experimental, temperature dependent excitation spectrum in the JILA TOP trap (filled circles) versus the HFB-Popov predictions for the $m = 0$ mode (top, labeled by "+") and the $m = 2$ mode (bottom, labeled by "x"). The solid curves are excitation frequencies for a zero-temperature condensate having the same number of condensate atoms as the experimental condensate in the finite- T cloud.




FIG. 2. A plot of the condensate fraction as a function of T' (solid curve) for the JILA TOP trap in which N is fixed at 2000 atoms. The same quantity is shown for the ideal gas in the thermodynamic limit (dotted curve) and for 2000 atoms (dashed curve) for comparison.




FIG. 3. HFB-Popov excitation frequencies (filled circles) for the $m = 0$ (top), $m = 2$ (middle), and the $m = 1$ modes (bottom) for a cold-atom cloud having $N = 2000$ atoms. Overlaid (solid lines) are the frequencies for a zero-temperature system with the same number N_0 of condensate atoms as in the finite-temperature system.